

Revised: 10/3/2016 at 7:37 Al

## Name:

|         | Take These Medications                              | At These Times  | Purpose   |
|---------|---|---|---|
|         | - Remind Me   |   |   |
| GENERIC | <b>Aspirin</b><br>81 mg Enteric coated tablet(s)    | take one tablet daily at<br>breakfast time  | Prevents blood clots  |
| GENERIC | Carvedilol<br>3.125 mg Tablet(s)                    | take one tablet at<br>breakfast and one tablet<br>at dinner time                                  | Controls blood pressure;<br>Heart medicine                                  |
| GENERIC | Furosemide<br>20 mg Tablet(s)                       | take one tablet at<br>breakfast time and one<br>tablet at dinner time                             | Water pill  |
| GENERIC | <b>Lisinopril</b><br>2.5 mg Tablet(s)               | take two tablets at<br>breakfast time   | Controls blood pressure;<br>Heart medicine                                  |
| 7       | <b>Xarelto®</b><br>(Rivaroxaban)<br>15 mg Tablet(s) | take one tablet at<br>breakfast time and one<br>tablet at dinner time.<br>Must be taken with food | Reduces the risk of stroke<br>in patients that have<br>irregular heart beat |
| GENERIC | Spironolactone<br>50 mg Tablet(s)                   | take one tablet at<br>bedtime which is at 10pm  | Water pill  |

The pharmacist may make a generic substitution for the medication shown in your Daily Schedule. The medication name, dosage strength, shape, color, and size may change as a result of this substitution. Please check dosage strength on your prescription bottle against the dosage strength shown on your daily schedule. Call your healthcare professional's office immediately if you receive medication that is different from what was prescribed or if you have questions about your medication. 1 teaspoon = 5mL, 1 tablespoon = 15mL

©2001-2016 McdActionPlan.com, LLC. All Rights Reserved.



DONATE Save a life, become an organ donor DonateLife.net

