



**DIABETES  
COLLABORATIVE  
REGISTRY®**

**PINNACLE Registry®**

Measure	Measure Title	Measure Description	Measure Type	NQS Domain	Meaningful Measure Area	Comparable PINN / DC ID
<a href="#"><u>QPP 326</u></a> (NQF 1525)	Atrial Fibrillation and Atrial Flutter: Chronic Anticoagulation Therapy	Percentage of patients aged 18 years and older with nonvalvular atrial fibrillation (AF) or atrial flutter who were prescribed warfarin OR another FDA-approved oral anticoagulant drug for the prevention of thromboembolism during the measurement period.	Process	Effective Clinical Care	Management of Chronic Conditions	PINN-161
<a href="#"><u>QPP 243</u></a> (NQF 0643)	Cardiac Rehabilitation Patient Referral from an Outpatient Setting	Percentage of patients evaluated in an outpatient setting who within the previous 12 months have experienced an acute myocardial infarction (MI), coronary artery bypass graft (CABG) surgery, a percutaneous coronary intervention (PCI), cardiac valve surgery, or cardiac transplantation, or who have chronic stable angina (CSA) and have not already participated in an early outpatient cardiac rehabilitation/secondary prevention (CR) program for the qualifying event/diagnosis who were referred to a CR program.	Process High Priority	Communication and Care Coordination	Preventive Care	PINN-108
<a href="#"><u>QPP 47</u></a> (NQF 0326)	Advance Care Plan	Percentage of patients aged 65 years and older who have an advance care plan or surrogate decision maker documented in the medical record or documentation in the medical record that an advance care plan was discussed but the patient did not wish or was not able to name a surrogate decision maker or provide an advance care plan.	Process High Priority	Communication and Care Coordination	Care is Personalized and Aligned with Patient's Goals	PINN-14 DC-10
<a href="#"><u>QPP 118</u></a> (NQF 0066)	Coronary Artery Disease: Angiotensin-Converting Enzyme (ACE) Inhibitor or Angiotensin Receptor Blocker (ARB) Therapy- Diabetes or Left Ventricular Systolic Dysfunction (LVEF < 40%)	Percentage of patients aged 18 years and older with a diagnosis of coronary artery disease seen within a 12-month period who also have diabetes OR a current or prior Left Ventricular Ejection Fraction (LVEF) < 40% who were prescribed ACE inhibitor or ARB therapy.	Process	Effective Clinical Care	Management of Chronic Conditions	PINN-107 DC-2
<a href="#"><u>QPP 6</u></a> (NQF 0067)	Coronary Artery Disease: Antiplatelet Therapy	Percentage of patients aged 18 years and older with a diagnosis of coronary artery disease (CAD) seen within a 12-month period who were prescribed aspirin or clopidogrel.	Process	Effective Clinical Care	Management of Chronic Conditions	PINN-105
<a href="#"><u>QPP 7</u></a> (NQF 0070)	Coronary Artery Disease (CAD): Beta-Blocker Therapy – Prior Myocardial Infarction (MI) or Left Ventricular Systolic Dysfunction (LVEF < 40%)	Percentage of patients aged ≥18 years with a diagnosis of coronary artery disease seen within a 12-month period who also have prior myocardial infarction or a current or prior LVEF <40% who were prescribed beta-blocker therapy.  <b>*2 performance rates*</b>	Process	Effective Clinical Care	Management of Chronic Conditions	PINN-106
<a href="#"><u>eCQM:CMS145v7</u></a>	Coronary Artery Disease (CAD): Beta-Blocker Therapy – Prior Myocardial Infarction (MI) or Left Ventricular Systolic Dysfunction (LVEF < 40%)	Percentage of patients aged ≥18 years with a diagnosis of coronary artery disease seen within a 12-month period who also have prior myocardial infarction or a current or prior LVEF <40% who were prescribed beta-blocker therapy.  <b>*2 performance rates*</b>	Process	Effective Clinical Care	Management of Chronic Conditions	PINN-106



**DIABETES  
COLLABORATIVE  
REGISTRY®**

**PINNACLE Registry®**

Measure	Measure Title	Measure Description	Measure Type	NQS Domain	Meaningful Measure Area	Comparable PINN / DC ID
<a href="#">QPP 1</a> (NQF 0059)	Diabetes: Hemoglobin A1c Poor Control	Percentage of patients 18-75 years of age with diabetes who had hemoglobin A1c > 9.0% during the measurement period.	Intermediate Outcome High Process Inverse	Effective Clinical Care	Management of Chronic Conditions	DC-1
<a href="#">eCQM:CMS122v7</a>	Diabetes: Hemoglobin A1c Poor Control	Percentage of patients 18-75 years of age with diabetes who had hemoglobin A1c > 9.0% during the measurement period (12 months from date of encounter).	Outcome Inverse	Effective Clinical Care	Management of Chronic Conditions	DC-1
<a href="#">QPP 119</a> (NQF 0062)	Diabetes: Medical Attention for Nephropathy	The percentage of patients 18-75 years of age with diabetes who had a nephropathy-screening test or evidence of nephropathy during the measurement period.	Process	Effective Clinical Care	Management of Chronic Conditions	DC-3
<a href="#">eCQM:CMS134v7</a>	Diabetes: Medical Attention for Nephropathy	The percentage of patients 18-75 years of age with diabetes who had a nephropathy-screening test or evidence of nephropathy during the measurement period.	Process	Effective Clinical Care	Management of Chronic Conditions	DC-3
<a href="#">ARCO 15</a>	Diabetes: Nutritional, Weight Loss Counseling, Reduction of Sedentary Behavior	Percentage of diabetic patients receiving nutritional counseling.	Process	Community / Population Health	Management of Chronic Conditions	
<a href="#">QPP 130</a> (NQF 0419)	Documentation of Current Medications in the Medical Record	Percentage of visits for patients aged 18 years and older for which the eligible clinician attests to documenting a list of current medications using all immediate resources available on the date of the encounter. This list must include ALL known prescriptions, over-the-counters, herbals, and vitamin/mineral/dietary (nutritional) supplements AND must contain the medications' name, dosage, frequency and route of administration.	Process High Priority	Patient Safety	Medication Management	
<a href="#">eCQM:CMS68v8</a>	Documentation of Current Medications in the Medical Record	Percentage of visits for patients aged 18 years and older for which the eligible clinicians attests to documenting a list of current medications using all immediate resources available on the date of the encounter. This list must include ALL known prescriptions, over-the-counters, herbals, and vitamin/mineral/dietary (nutritional) supplements AND must contain the medications' name, dosage, frequency and route of administration.	Process	Patient Safety	Medication Management	
<a href="#">QPP 8</a> (NQF 0083)	Heart Failure (HF): Beta-Blocker Therapy for Left Ventricular Systolic Dysfunction (LVSD)	Percentage of patients aged 18 years and older with a diagnosis of heart failure (HF) with a current or prior left ventricular ejection fraction (LVEF) < 40% who were prescribed beta-blocker therapy either within a 12-month period when seen in the outpatient setting OR at each hospital discharge.	Process	Effective Clinical Care	Management of Chronic Conditions	PINN-144



**DIABETES  
COLLABORATIVE  
REGISTRY®**

**PINNACLE Registry®**

Measure	Measure Title	Measure Description	Measure Type	NQS Domain	Meaningful Measure Area	Comparable PINN / DC ID
<a href="#">eCQM:CMS144v7</a>	Heart Failure: Beta-Blocker Therapy for Left Ventricular Systolic Dysfunction LVSD	Percentage of patients aged 18 years and older with a diagnosis of heart failure (HF) with a current or prior left ventricular ejection fraction (LVEF) < 40% who were prescribed beta-blocker therapy either within a 12-month period when seen in the outpatient setting OR at each hospital discharge.	Process	Effective Clinical Care	Management of Chronic Conditions	PINN-144
<a href="#">QPP 5 (NQF 0081)</a>	Heart Failure: Angiotensin-Converting Enzyme (ACE) Inhibitor or Angiotensin Receptor Blocker (ARB) Therapy for Left Ventricular Systolic Dysfunction (LVSD)	Percentage of patients aged ≥18 years and older with a diagnosis of HF with a current or prior LVEF of <40% who were prescribed ACE inhibitor or ARB therapy either within a 12-month period when seen in the outpatient setting OR at each hospital discharge.	Process	Effective Clinical Care	Management of Chronic Conditions	PINN-145
<a href="#">eCQM:CMS135v7</a>	Heart Failure: Angiotensin-Converting Enzyme (ACE) Inhibitor or Angiotensin Receptor Blocker (ARB) Therapy for Left Ventricular Systolic Dysfunction (LVSD)	Percentage of patients aged 18 years and older with a diagnosis of heart failure (HF) with a current or prior left ventricular ejection fraction (LVEF) < 40% who were prescribed ACE inhibitor or ARB therapy either within a 12-month period when seen in the outpatient setting OR at each hospital discharge.	Process	Effective Clinical Care	Management of Chronic Conditions	PINN-145
<a href="#">ACCPin8</a>	Hypertension: Blood Pressure Control for Stage 1 or 2 Hypertensive Patients	Proportion of patients with hypertension who had adequately controlled blood pressure.	Process	Effective Clinical Care	Management of Chronic Conditions	PINN-123
<a href="#">ACCPin9</a>	Hypertension: Blood Pressure Treatment and Control for High Risk Patients	Proportion of patients with both hypertension and a ≥ 10% CVD risk OR high-risk diagnosis (i.e. ASCVD, chronic kidney disease, diabetes) who were prescribed antihypertensive medication or who had adequately controlled blood pressure.	Process	Effective Clinical Care	Management of Chronic Conditions	PINN-121
<a href="#">ACCPin10</a>	Hypertension: Blood Pressure Treatment and Control for Stage 2 Hypertensive Patients	Proportion of patients with hypertension who were prescribed antihypertensive medications or who had adequately controlled blood pressure.	Process	Effective Clinical Care	Management of Chronic Conditions	PINN-122
<a href="#">ACCPin7</a>	Peripheral Artery Disease: Treatment of Blood Cholesterol to Reduce Atherosclerotic Cardiovascular Risk	Percentage of Patients 18-75 years of age with PAD who were offered moderate-to-high intensity statin.	Process	Effective Clinical Care	Management of Chronic Conditions	PINN-170 DC-7
<a href="#">QPP 441</a>	Ischemic Vascular Disease (IVD) All or None Outcome Measure (Optimal Control)	The IVD All-or-None Measure is one outcome measure (optimal control). The measure contains four goals. All four goals within a measure must be reached in order to meet that measure. The numerator for the all-or-none measure should be collected from the organization's total IVD denominator. All-or-None Outcome Measure (Optimal Control) - Using the IVD denominator optimal results include: <ul style="list-style-type: none"> <li>• Most recent blood pressure (BP) measurement is less than or equal to 140/90 mm Hg -- AND</li> <li>• Most recent tobacco status is Tobacco Free -- AND</li> <li>• Daily Aspirin or Other Antiplatelet Unless Contraindicated -- AND</li> <li>• Statin Use Unless Contraindicated</li> </ul>	Intermediate Outcome High Priority	Effective Clinical Care	Management of Chronic Conditions	



Measure	Measure Title	Measure Description	Measure Type	NQS Domain	Meaningful Measure Area	Comparable PINN / DC ID
<a href="#">QPP 128</a> (NQF 0421)	Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan	Percentage of patients aged 18 years and older with a BMI documented during the current encounter or during the previous twelve months AND with a BMI outside of normal parameters, a follow-up plan is documented during the encounter or during the previous twelve months of the current encounter.	Process	Community/Population Health	Preventive Care	
<a href="#">eCQM:CMS69v7</a>	Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan	Patients with a documented BMI during the encounter or during the previous twelve months, AND when the BMI is outside of normal parameters, a follow-up plan is documented during the encounter or during the previous twelve months of the current encounter.	Process	Community/Population Health	Preventive Care	
<a href="#">QPP 226</a> (NQF 0028)	Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention (AMA-PCPI)	Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 24 months AND who received tobacco cessation intervention if identified as a tobacco user.  <b>*3 performance rates*</b>	Process	Community/Population Health	Prevention and Treatment of Opioid and Substance Use Disorders	PINN-104 DC-4
<a href="#">eCQM:CMS138v7</a>	Tobacco Use: Screening and Cessation Intervention	Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 24 months AND who received tobacco cessation intervention if identified as a tobacco user.  <b>*3 performance rates*</b>	Process	Community/Population Health	Prevention and Treatment of Opioid and Substance Use Disorders	PINN-104 DC-4

### 2019 QPP Requirements

**Individual Provider Reporting:** Eligible providers must report at least six measures, including one outcome measure. If an outcome measure is not available that is applicable to your specialty or practice, choose a high priority measure. Measures must be calculated with at least 60 percent of the providers patients (all-payer) included in the denominator.

**Group Practice Reporting (GPRO):** A group is defined as a single tax identification number (TIN) with two or more clinicians (at least one clinician within the group must be MIPS eligible) as identified by their national provider identifier (NPI), who have reassigned their Medicare billing rights to a single TIN must report at least six measures, including one outcome measure. If an outcome measure is not available that is applicable to your specialty or practice, choose a high priority measure. Measures must be calculated with at least 60 percent of the providers patients (all-payer) included in the denominator.

**Measure specifications for QPP measures are available where hyperlinked. All measures are proportional, traditional (unless indicated differently under *Measure Type*) and NOT risk adjusted.**